

$$2) \quad F_{z\bar{z}} \bar{z} \gamma^{z\bar{z}} \Sigma + \mathcal{D} = F_{z\bar{z}} \Sigma^{z\bar{z}} + \mathcal{D} = 0$$

along Riemann surface

→ solved by $F_{z\bar{z}} = \text{const. } \Sigma_{z\bar{z}}$

allowed constants are quantized as:

$$\frac{1}{24} \int_{\Sigma} F = c_1(F) \in \mathbb{Z}$$

can add $g-1$ additional flat connections

→ for each $u \in \mathbb{Z}$ and each $u(1) \subset G$
get g complex moduli

More general possibility:

fix L abelian subgroup of G

→ commutant: $L \times G' \subset G$,

fix $c_1(L) \in \mathbb{Z}^{\dim(L)} \rightarrow g \cdot \dim(L)$ abelian moduli

+ $(g-1) \dim(G')$
non-abelian moduli

expected dimension of 4d $\mathcal{N}=1$ conformal manifold \mathcal{M}_g :

$$\dim \mathcal{M}_g = (3g-3) + g \cdot \dim(L) + (g-1) \dim(G')$$

Example 1:

6d (2,0) SCFT with $SO(5)_R$ R-symmetry

$$\text{we have } su(2)_L \times su(2)_R = SO(4) \subset SO(5)$$

$$\cup$$
$$SO(2)_R \text{ of 4d } \mathcal{N}=1$$

$$t_1 + t_2$$
$$SO(2) \times SO(2) \subset SO(4)$$

→ flavor symmetry $G = su(2)_L$

1) turn on flat $su(2)_L$ bundle on Σ_g :

$$\dim \mathcal{M}_g = (3g-3) + (g-1) \cdot 3 = 6g-6$$

2) Choose abelian subgroup $L \subset G : u(1) \subset su(2)_L$

→ turn on flux characterized
by integer $c(L) = n$

$$\rightarrow \dim \mathcal{M}_g = (3g-3) + g = 4g-3$$

Example 2:

consider E-string $\mathcal{N}=(1,0)$ SCFT in 6d

→ global symmetry: $G = E_8$

1) choose $G' = G = E_8$ → moduli space of flat
 E_8 bundles on Σ_g

2) or abelian subgroup $L \subset E_8$

with $\dim L = 1, \dots, 8$

→ numerous possibilities for each dimension

Contrast this with $\mathcal{N}=(2,2)$ SCFT


→ no global symmetries: $G=0$

→ unique choice

Adding punctures:

S punctures → positions add S complex moduli

semi-infinite cylinder



↓ reduction on circle
+ of holonomy for G

5d theory

+ mass-parameters

→ G is broken to $P \subset G$

inequivalent choices: G/P

→ each puncture adds complex dim

$\frac{1}{2} \dim(G/P)$ moduli

holonomies in the bulk: $P^{\max} = G^{\max} \cap P$

where $G^{\max} = L \times G'$

→ dimension of 4d conformal manifold:

$$\begin{aligned} \dim \mathcal{M}_g &= (3g-3+s) + (g-1) \dim(G^{\max}) + \dim(L) \\ &\quad + \frac{1}{2} \sum_i \dim\left(\frac{G^{\max}}{P_i^{\max}}\right) \\ &= 3g-3+s + \left(g-1 + \frac{s}{2}\right) \dim(G^{\max}) \\ &\quad + \dim(L) - \frac{1}{2} \sum_i \dim(P_i^{\max}) \end{aligned}$$

M5 branes probing ADE compactified on Σ_g

N M5 branes probing \mathbb{C}^2/Γ_k

→ resulting theory has $G = K_L \times K_R$
global symmetry

focus on $K = \text{SU}(k)$

For $N=1$: 6d SCFT is hyper in $(k, \bar{k})_{+1}$

representation of $\text{SU}(k)_L \times \text{SU}(k)_R \times \text{U}(1)$

can turn on non-abelian bundles for
 $\text{SU}(k)_L \times \text{SU}(k)_R$ along cycles of Σ_g
and abelian bundles from $\text{U}(1)$